

A Note on the Andrica Conjecture

Marek Wolf

e-mail: primes7@o2.pl

Abstract

We derive heuristically the approximate formula for the difference $\sqrt{p_{n+1}} - \sqrt{p_n}$, where p_n is the n -th prime. We find perfect agreement between this formula and the available data from the list of maximal gaps between consecutive primes.

1 Introduction

The Andrica conjecture [1] (see also [7, p.21] and [12, p. 191]) states that the inequality:

$$A_n \equiv \sqrt{p_{n+1}} - \sqrt{p_n} < 1 \quad (1)$$

where p_n is the n -th prime number, holds for all n . Despite its simplicity it remains unproved. In the Table I we give a few first values of A_n and in Table II the values of A_n are sorted in descending order.

We have

$$\sqrt{p_{n+1}} - \sqrt{p_n} = \frac{p_{n+1} - p_n}{\sqrt{p_{n+1}} + \sqrt{p_n}} < \frac{d_n}{2\sqrt{p_n}} \quad (2)$$

From this we see that the growth rate of the form $d_n = \mathcal{O}(p_n^\theta)$ with $\theta < 1/2$ will suffice for the proof of (1). Unfortunately all values of θ proved in the past are larger than $1/2$. A few results with θ closest to $1/2$ are: M. Huxley: $\theta > 7/12$ [8], the result of C.J. Mozzochi [10] $\theta = \frac{1051}{1920}$, S. Lou and Q. Yao obtained $\theta = 6/11$ [9] and recently R.C. Baker G. Harman and J. Pintz [2] have improved it to $\theta = 21/40$

what remains currently the best unconditional result. For a review of results on θ see [11]. The best estimation for d_n obtained by H. Cramer under the assumption of the Riemann Hypothesis [3]

$$d_n = \mathcal{O}(\sqrt{p_n} \log(p_n)) \quad (3)$$

also does not suffice to prove the Andrica conjecture.

TABLE I

p_n	p_{n+1}	d_n	$\sqrt{p_{n+1}} - \sqrt{p_n}$	p_n	p_{n+1}	d_n	$\sqrt{p_{n+1}} - \sqrt{p_n}$
2	3	1	0.317837245	41	43	2	0.154314287
3	5	2	0.504017170	43	47	4	0.298216076
5	7	2	0.409683334	47	53	6	0.424455289
7	11	4	0.670873479	53	59	6	0.401035859
11	13	2	0.288926485	59	61	2	0.129103928
13	17	4	0.517554350	61	67	6	0.375103096
17	19	2	0.235793318	67	71	4	0.240797001
19	23	4	0.436932580	71	73	2	0.117853972
23	29	6	0.589333284	73	79	6	0.344190672
29	31	2	0.182599556	79	83	4	0.222239162
31	37	6	0.514998167	83	89	6	0.323547553
37	41	4	0.320361707	89	97	8	0.414876670
41	43	2	0.154314287	97	101	4	0.201017819
43	47	4	0.298216076	101	103	2	0.099015944
47	53	6	0.424455289	103	107	4	0.195188868
53	59	6	0.401035859	107	109	2	0.096226076
59	61	2	0.129103928	109	113	4	0.189839304

For twins primes $p_{n+1} = p_n + 2$ there is no problem with (1) and in general for short gaps $d_n = p_{n+1} - p_n$ between consecutive primes the inequality (1) will be satisfied. The Andrica conjecture can be violated only by extremely large gaps

between consecutive primes. Let $G(x)$ denote the largest gap between consecutive primes smaller than x :

$$G(x) = \max_{p_n, p_{n-1} < x} (p_n - p_{n-1}). \quad (4)$$

Let us denote the pair of primes $< x$ comprising the largest gap $G(x)$ by $p_{L+1}(x)$ and $p_L(x)$, hence we have

$$G(x) = p_{L+1}(x) - p_L(x). \quad (5)$$

Thus we will concentrate on the values of the difference appearing in (1) corresponding to the largest gaps and let us introduce the function:

$$R(x) = \sqrt{p_{L+1}(x)} - \sqrt{p_L(x)} \quad (6)$$

Then we have:

$$A_n \leq R(p_n). \quad (7)$$

The largest values of A_n will be reached at the largest gaps $G(x)$ between consecutive primes below a given bound x . In [15], [14] we have given the heuristic arguments that $G(x)$ can be expressed directly by $\pi(x)$ — the number of primes $< x$:

$$G(x) \sim \frac{x}{\pi(x)} (2 \log \pi(x) - \log(x) + c'), \quad (8)$$

where c' is expressed by the twin constant C_2 :

$$c' = \log(C_2) = 0.27787688 \dots, \quad C_2 \equiv 2 \prod_{p>2} \left(1 - \frac{1}{(p-1)^2} \right) = 1.32032363169 \dots \quad (9)$$

For the Gauss approximation $\pi(x) \sim x/\log(x)$ the following dependence follows:

$$G(x) \sim \log(x)(\log(x) - 2 \log \log(x) + \log(c')) \quad (10)$$

and for large x it passes into the Cramer [4] conjecture:

$$G(x) \sim \log^2(x). \quad (11)$$

TABLE II

n	p_n	p_{n+1}	d_n	$\sqrt{p_{n+1}} - \sqrt{p_n}$
4	7	11	4	0.6708735
30	113	127	14	0.6392819
9	23	29	6	0.5893333
6	13	17	4	0.5175544
11	31	37	6	0.5149982
2	3	5	2	0.5040172
8	19	23	4	0.4369326
15	47	53	6	0.4244553
46	199	211	12	0.4191031
34	139	149	10	0.4167295

A. Granville argued [6] that the actual $G(x)$ can be larger than that given by (11), namely he claims that there are infinitely many pairs of primes p_n, p_{n+1} for which:

$$p_{n+1} - p_n = G(p_n) > 2e^{-\gamma} \log^2(p_n) = 1.12292 \dots \log^2(p_n). \quad (12)$$

For a given gap d the largest value of the difference $\sqrt{p+d} - \sqrt{p}$ will appear at the first appearance of this gap: each next pair $(p', p' + d)$ of consecutive primes separated by d will produce smaller difference (see (2)):

$$\sqrt{p'+d} - \sqrt{p'} < \sqrt{p+d} - \sqrt{p}. \quad (13)$$

Hence we have to focus our attention on the first occurrences of gaps. In [16] we have given heuristic arguments that the gap d should appear for the first time after the prime $p_f(d)$ given by

$$p_f(d) \sim \sqrt{d} e^{\sqrt{d}}. \quad (14)$$

We calculate

$$\begin{aligned}\sqrt{p_f(d) + d} - \sqrt{p_f(d)} &= \sqrt{\sqrt{d}e^{\sqrt{d}} + d} - \sqrt{\sqrt{d}e^{\sqrt{d}}} = \\ &\sqrt{\sqrt{d}e^{\sqrt{d}}} \left(\sqrt{1 + \frac{d}{\sqrt{d}e^{\sqrt{d}}}} - 1 \right) = \frac{1}{2}d^{\frac{3}{4}}e^{-\frac{1}{2}\sqrt{d}} + \dots\end{aligned}\quad (15)$$

Substituting here for d the maximal gap $G(x)$ given by (8) we obtain the approximate formula for $R(x)$:

$$R(x) = \frac{1}{2}G(x)^{3/4}e^{-\frac{1}{2}\sqrt{G(x)}} + \text{error term}.\quad (16)$$

The comparison with real data is given in Figure 1. The lists of known maximal gaps between consecutive primes can be found at <http://www.trnicely.net> and <http://www.ieeta.pt/~tos/gaps.html>. The largest known gap 1476 between consecutive primes follows the prime $1425172824437699411 = 1.42 \dots \times 10^{18}$.

The maximum of the function $\frac{1}{2}x^{\frac{3}{4}}e^{-\frac{1}{2}\sqrt{x}}$ is reached at $x = 9$ and has the value 0.579709161122. The maximal value of A_n is 0.6708735... for $d = 4$ and second value is 0.6392819... for $d = 14$. Let us remark that $d = 9$ is exactly in the middle between 4 and 14.

Because in (16) $R(x)$ contains exponential of $\sqrt{G(x)}$ it is very sensitive to the form of $G(x)$. The substitution $G(x) = \log^2(x)$ leads to the form:

$$R(x) = \frac{\log^{3/2}(x)}{2\sqrt{x}}.\quad (17)$$

This form of $R(x)$ is plotted in Fig.2 in red. In [13] D. Shanks has given for $p_f(d)$ the expression

$$p_f(d) \sim e^{\sqrt{d}}.\quad (18)$$

This leads to the expression

$$\sqrt{p_f(d) + d} - \sqrt{p_f(d)} = \frac{1}{2}de^{-\frac{1}{2}\sqrt{d}}\quad (19)$$

instead of (15). Substitution here for d the form (10) leads to the curve plotted in Fig.2 in green.

Finally let us remark, that from the above analysis it follows, that

$$\lim_{n \rightarrow \infty} \sqrt{p_{n+1}} - \sqrt{p_n} = 0\quad (20)$$

The above limit was mentioned on p. 61 in [5] as a difficult problem (yet unsolved).




Fig.1 The plot of $R(x)$ and approximation to it given by (main-formula). The are 75 maximal gaps available currently and hence there are 75 circles in the plot of $R(x)$.




Fig.1 The plot of $R(x)$ and approximation to it given by $\frac{\log^{3/2}(x)}{2\sqrt{x}}$ (red) and approximation of $R(x)$ obtained from the Shanks conjecture for $p_f(d)$ (green).

References

- [1] D. Andrica. Note on a conjecture in prime number theory. *Studia Univ. Babes-Bolyai Math.*, 31:44–48, 1986.
- [2] R. C. Baker, G. Harman, and P. J. The difference between consecutive primes, II. *Proc. London Math. Soc.*, 83:532562, 2001.
- [3] H. Cramer. Some theorems concerning prime numbers. *Arkiv f. Math. Astr. Fys.*, 15:1–33, 1920.
- [4] H. Cramer. On the order of magnitude of difference between consecutive prime numbers. *Acta Arith.*, II:23–46, 1937.
- [5] S. W. Golomb. Problem E2506: Limits of differences of square roots. *Amer. Math. Monthly*, 83:60–61, 1976.
- [6] A. Granville. Harald Cramer and the distribution of prime numbers. *Scandinavian Actuarial J.*, 1:12–28, 1995.

- [7] R. K. Guy. *Unsolved Problems in Number Theory*. Springer-Verlag, 2nd ed. New York, 1994.
- [8] M. Huxley. An application of the Fouvry-Iwaniec theorem. *Acta Arithmetica*, 43:441–443, 1984.
- [9] S. Lou and Q. Yao. A Chebyshev’s type of prime number theorem in a short interval. ii. *Hardy-Ramanujan J.*, 15:1–33, 1992.
- [10] C. Mozzochi. On the difference between consecutive primes. *Journal Number Theory*, 24:181–187, 1986.
- [11] J. Pintz. Landaus problems on primes. *Journal de thorie des nombres de Bordeaux*, 21:357–404, 2009.
- [12] P. Ribenboim. *The Little Book of Big Primes*. 2ed., Springer, 2004.
- [13] D. Shanks. On maximal gaps between successive primes. *Mathematics of Computation*, 18:646–651, 1964.
- [14] M. Wolf. Some conjectures on the gaps between consecutive primes, 1995. preprint IFTUWr 894//95, September 1995, available from <http://www.ift.uni.wroc.pl/~mwolf/conjectures.ps.gz>.
- [15] M. Wolf. Unexpected regularities in the distribution of prime numbers. In P. et al, editor, *8th Joint EPS-APS Int.Conf. Physics Computing’96, Krakw, 1996*, pages 361–367, 1996.
- [16] M. Wolf. First occurence of a given gap between consecutive primes, 1997. preprint IFTUWr 911//97, April 1997, available from <http://www.ift.uni.wroc.pl/~mwolf/firstocc.pdf>.